

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

at D224
W2W37

80



Washington Basin Outlook Report

May 1, 1998



Basin Outlook Reports

and

Federal - State - Private

Cooperative Snow Surveys

For more water supply and resource management information, contact:

Local Natural Resources Conservation Service Field Office

or

Scott Pattee
Water Supply Specialist
Natural Resources Conservation Service
2021 E. College Way, Suite 214
Mt. Vernon, WA 98273-2873
(360) 428-7684

or

Chris Bieker
Public Affairs Specialist
Natural Resources Conservation Service
316 W. Boone Ave., Suite 450
Spokane, WA 99201-2348
(509) 323-2912

How forecasts are made

Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snow courses and automated SNOTEL sites, along with precipitation, antecedent streamflow, and indices of the El Niño / Southern Oscillation are used in computerized statistical and simulation models to prepare runoff forecasts. These forecasts are coordinated between hydrologists in the Natural Resources Conservation Service and the National Weather Service. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less water.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all of its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require an alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue SW, Washington DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an Equal Opportunity provider and employer.

Washington Water Supply Outlook

May 1998

General Outlook

Predictably, snowpack levels dropped dramatically over most of the state. Above average temperatures and below average precipitation was the basic theme for the month. Streamflow forecasts for some Washington streams took another dive however a select few actually increased. April streamflows varied across the state, but on average were near normal. Reservoir storage varied greatly from basin to basin. Temperatures for the month were 1 to 3 degrees above normal.

Snowpack

The May 1 statewide SNOTEL readings showed 102% of average snowpack. By May 4 the average had dropped to 91%, due to unseasonably high temperatures. Snowpack varied from 28% of average in the Tolt River Basin to as high as 436% in the Conconully Lake Basin. Westside averages, from SNOTEL and May 1 snow surveys, included the North Puget Sound river basins with 78% of average, the Olympic Peninsula basins with 104%, and the Lewis-Cowlitz basins with 117% of average. Snowpack along the east slopes of the Cascade Mountains included the Yakima area with 124% of average, and the Wenatchee area with 161%. Snowpack in the Spokane River Basin remained below average at 50%, and the Pend Oreille River Basin (including Canadian data) retained only 65% of average snowpack. Maximum snow cover in the state was at Paradise SNOTEL near Mount Rainier. It had a water content of 70 inches. This site would normally have 61.8 inches of water content on May 1. The highest average in the state was Salmon Meadows SNOTEL in the Conconully Lake Basin with 436% of average.

BASIN	PERCENT OF LAST YEAR	PERCENT OF AVERAGE
Spokane	32	50
Newman Lake	34	52
Colville	N/A	N/A
Pend Oreille	40	65
Okanogan	60	82
Similkameen	44	57
Methow	62	95
Chelan	63	100
Wenatchee	48	90
Steilacoom Creek	58	144
Yakima	54	106
Ahtanum Creek	74	143
Walla Walla	30	66
Cowlitz	64	106
Lewis	56	129
White	76	123
Green	34	78
Cedar	24	48
Snoqualmie	41	78
Skykomish	32	65
Skagit	56	85
Baker	70	93
Nooksack	42	56
Olympic Peninsula	98	104

Precipitation

During the month of April, the National Weather Service and Natural Resources Conservation Service climate stations showed considerable variations in precipitation across Washington. The highest percent of average in the state was at Winthrop and Mazama, both in the Methow River Valley. Both climate stations reported 194% of average for a total of 1.49 and 1.96 inches respectfully. The April average for Winthrop was 0.77 inches and for Mazama it was 1.01 inches. Averages for the water year varied from 110% of average on the Olympic Peninsula to 80% in the Walla Walla River Basin. The highest individual site average for the water year was 160% of average at Trough SNOTEL site near Wenatchee.

RIVER BASIN	APRIL PERCENT OF AVERAGE	WATER YEAR PERCENT OF AVERAGE
Spokane	61	83
Colville-Pend Oreille	59	90
Okanogan-Methow	93	105
Wenatchee-Chelan	64	104
Yakima	58	102
Walla Walla	63	80
Cowlitz-Lewis	50	105
White-Green	50	93
Central Puget Sound	47	92
North Puget Sound	47	90
Olympic Peninsula	65	110

Reservoir

Storage levels are beginning to stabilize with the start of spring runoff and the irrigation season. However managers are still trying to balance anticipated spring runoff with summer usage and carry-over requirements. Reservoir storage in the Yakima Basin was 958,400 acre feet, or 122% of average. Storage at other reservoirs included the Okanogan reservoirs with 149% of average for May 1. Storage levels at Roosevelt and Banks Lakes was not available at publication time. The power generation reservoirs included the following: Coeur d'Alene Lake, 181,500 acre feet, or 74% of average and 76% of capacity; Chelan Lake, 312,800 acre feet, 70% of average and 46% of capacity; and the Skagit River reservoirs at 112% of average and 51% of capacity.

BASIN	PERCENT OF CAPACITY	PERCENT OF AVERAGE
Spokane	76	74
Colville-Pend Oreille	N/A	N/A
Okanogan-Methow	101	149
Wenatchee-Chelan	46	70
Yakima	90	122
North Puget Sound	51	112

Streamflow

Early season snow melt has brought forecasts for many Washington streams down even further this month. A few select streams are forecast slightly higher. Forecasts varied from 122% of average for Salmon Creek near Conconully, to 50% of average for the Spokane River near Post Falls. May forecasts for some Western Washington streams include: Cedar River near Cedar Falls, 77% of average; Green River, 68%; and the Dungeness River, 97%. Some Eastern Washington streams include the Yakima River near Parker, 88% of average; the Wenatchee River at Plain, 93%; and the Colville River at Kettle Falls, 93%. Volumetric forecasts are developed using current, historic, and average snowpack, precipitation and streamflow data collected and coordinated by organizations cooperating with NRCS. A beneficial fact sheet, "Interpreting Streamflow Forecasts," is available on the World Wide Web at <http://www.wcc.nrcs.usda.gov/factpub/factpub.html>

BASIN	PERCENT OF AVERAGE MOST PROBABLE FORECAST (50 PERCENT CHANCE OF EXCEEDENCE)
Spokane	50-57
Colville-Pend Oreille	59-105
Okanogan-Methow	73-122
Wenatchee-Chelan	93-101
Yakima	85-111
Walla Walla	77-93
Cowlitz-Lewis	89-96
Green River	68
Central Puget Sound	73-90
North Puget Sound	85-90
Olympic Peninsula	94-97

Streamflows reported for April continued to vary from well above to well below average. The Kettle River at Laurier, had the highest flows at 172% of average; and the Cowlitz River below Mayfield Dam, with 67% of average, had the lowest flows in the state. Other streamflows were the following percentage of average: the Priest River, 109%; the Columbia at the International Boundary, 110%; the Spokane River at Spokane, 74%; the Columbia below Rock Island Dam, 101%; the Cle Elum River near Roslyn, 98%; and the Snake River below Ice Harbor Dam, 95%.


STREAM	PERCENT OF AVERAGE APRIL STREAMFLOWS
Pend Oreille Below Box Canyon	98
Kettle at Laurier	172
Columbia at Birchbank	110
Spokane at Long Lake	79
Similkameen at Nighthawk	81
Okanogan at Tonasket	125
Methow at Pateros	133
Chelan at Chelan	108
Wenatchee at Pashastin	100
Yakima at Cle Elum	91
Yakima at Parker	100
Naches at Naches	111
Yakima at Kiona	103
Grande Ronde at Troy	77
Snake below Lower Granite Dam	100
SF Walla Walla near Milton Freewater	115
Columbia at The Dalles	101
Lewis at Ariel	68
Cowlitz below Mayfield Dam	67
Skagit at Concrete	79

For more information contact your local Natural Resources Conservation Service office.

**B A S I N S U M M A R Y O F
S N O W C O U R S E D A T A**


MAY 1998

SNOW COURSE	EL E V A T I O N	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 1961-90	SNOW COURSE	EL E V A T I O N	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 1961-90		
ALPINE MEADOWS PILL	3500	5/01/98	---	47.2S	72.6	--	MEADOWS CABIN	1900	4/29/98	0	.0	4.5	1.1		
AMBROSE	6480	4/28/98	22	9.4	22.4	12.1	MEADOWS PASS	PILLOW	3240	5/01/98	---	4.9S	35.0	21.0	
ASLEY DIVIDE	4820	4/28/98	0	.0	7.8	1.0	MERRITT		2140	5/01/98	0	.0	--	3.6	
BADGER PASS PILLOW	6900	5/01/98	---	20.6	49.1	37.8	MICA CREEK	PILLOW	4750	5/01/98	---	6.5	40.4	--	
BAREE CREEK	5500	4/28/98	64	31.2	61.8	43.0	MINERAL CREEK		4000	5/02/98	6	3.8	24.4	11.2	
BAREE MIDWAY	4600	4/28/98	46	20.8	53.0	29.4	MOOSE CREEK	PILLOW	6200	5/01/98	---	10.5	30.1	14.6	
BAREE TRAIL	3800	4/28/98	0	.0	13.8	1.3	MORRISSEY RIDGE CAN.		6100	5/01/98	---	18.1	41.8	28.7	
BARKER LAKES PILLOW	8250	5/01/98	---	12.7	23.8	16.0	MORSE LAKE	PILLOW	5400	5/01/98	---	68.3S	89.7	44.4	
BASIN CREEK PILLOW	7180	2/01/98	---	12.2	14.2	10.0	MOSES MTN	PILLOW	4800	5/01/98	---	12.2S	13.7	7.3	
BASSOO PEAK	5150	4/28/98	0	.0	11.4	5.7	MOSQUITO RDG	PILLOW	5200	5/01/98	---	20.6	58.1	34.7	
BEAVER CREEK TRAIL	2200	4/30/98	0	.0	16.8	4.1	MOULTON RESERVOIR		6850	4/24/98	0	.0	11.2	3.2	
BEAVER PASS	3680	4/30/98	47	22.4	42.3	28.1	MOUNT CRAG	PILLOW	4050	5/01/98	---	35.2S	28.4	22.4	
BERNE-MILL CREEK (d)	3170	5/01/98	44	20.1	38.3	20.8	MT. GARDNER	PILLOW	2860	5/01/98	---	.0S	18.8	10.8	
BIG CREEK	6750	4/27/98	95	36.8	70.1	49.8	N.F. ELK CR	PILLOW	6250	5/01/98	---	4.9	16.4	9.6	
BLACK MOUNTAIN	7750	4/29/98	46	16.6	23.2	17.8	NEVADA CREEK	PILLOW	6480	5/01/98	---	9.4	22.2	12.5	
BLACK PINE PILLOW	7100	5/01/98	---	7.2	18.5	12.0	NEW HOZOMEEN LAKE		2800	4/28/98	0	.0	13.2	4.5	
BLEWETT PASS#2PILLOW	4270	5/01/98	---	.5S	15.4	4.9	NEZ PERCE CMP	PILLOW	5650	5/01/98	---	8.0	21.4	11.7	
BLUE LAKE	5900	5/01/98	18	7.9	32.8	23.9	NEZ PERCE PASS		6570	4/28/98	28	12.4	22.6	15.6	
BROWN TOP	AM 6000	4/30/98	114	53.8	77.4	61.7	NOISY BASIN	PILLOW	6040	5/01/98	---	34.7	72.8	44.0	
BRUSH CREEK TIMBER	5000	4/28/98	0	.0	6.8	6.0	NORTH FORK JOCKO		6330	4/27/98	66	29.8	66.0	44.6	
BULL MOUNTAIN	6600	4/30/98	0	.0	11.4	3.1	OLALLIE MDWS	PILLOW	3960	5/01/98	---	49.9S	97.9	51.0	
BUMPING LAKE	3450	4/29/98	18	8.1	--	OLALLIE MEADOWS		3630	5/01/98	---	58.1E	--	43.5		
BUMPING LAKE (NEW)	3400	4/29/98	18	8.1	26.8	10.9	OPHIR PARK		7150	5/03/98	18	6.2	22.2	17.4	
BUMPING RIDGE PILLOW	4600	5/01/98	---	25.6S	48.9	18.9	PARADISE PARK	PILLOW	5500	5/01/98	---	70.0S	120.0	61.8	
BUNCHGRASS MDWPILLOW	5000	5/01/98	---	25.2	48.2	26.9	PARK CK RIDGE	PILLOW	4600	5/01/98	---	33.4S	69.6	33.6	
CAYUSE PASS	5300	5/01/98	---	96.0E	113.5	88.1	PETERSON MDW	PILLOW	7200	4/30/98	---	12.2	17.6	11.3	
CHESSMAN RESERVOIR	6200	4/29/98	0	.0	1.0	PIGTAIL PEAK	PILLOW	5900	5/01/98	---	50.4S	101.9	47.7		
CHICKEN CREEK	4060	4/27/98	0	.0	19.9	3.6	PIKE CREEK	PILLOW	5930	5/01/98	---	12.1	40.2	27.8	
CHIWAUKUN G.S.	2500	5/01/98	0	.0	--	PIPESTONE PASS		7200	4/29/98	21	7.4	10.0	5.0		
COMBINATION PILLOW	5600	5/01/98	---	.0	4.1	3.2	POPE RIDGE	PILLOW	3540	5/01/98	---	9.7S	23.5	1.6	
COPPER BOTTOM PILLOW	5200	5/01/98	---	.0	13.0	8.1	POTATO HILL	PILLOW	4500	5/01/98	---	19.0S	34.9	17.0	
COPPER MOUNTAIN	7700	4/28/98	32	10.8	17.8	QUARTZ PEAK	PILLOW	4700	5/01/98	---	9.7	28.4	18.6		
CORRAL PASS	PILLOW	6000	5/01/98	---	35.1S	59.5	29.5	ROUND TOP MTN		4020	4/29/98	0	.0	--	--
COTTONWOOD CREEK	6400	4/29/98	23	8.5	12.6	7.6	RAGGED RIDGE		3330	4/29/98	0	.0	--	--	
COUGAR MTN.	PILLOW	3200	5/01/98	---	.6S	30.2	9.3	RAINY PASS	PILLOW	4780	5/01/98	---	29.0S	54.7	36.8
COX VALLEY	4500	4/26/98	82	37.9	49.0	39.1	REX RIVER	PILLOW	1900	5/01/98	---	14.2S	43.7	23.1	
COYOTE HILL	4200	4/29/98	0	.0	12.1	3.0	ROCKER PEAK	PILLOW	8000	5/01/98	---	15.1	20.6	17.7	
DALY CREEK PILLOW	5780	5/01/98	---	6.5	16.6	5.8	SADDLE MTN	PILLOW	7900	5/01/98	---	22.6	42.9	27.6	
DEER PARK	5200	4/28/98	39	17.5	13.7	58.7	SALMON MDWS	PILLOW	4500	5/01/98	---	4.8S	9.0	1.1	
DEVILS PARK	5900	4/28/98	76	35.7	59.4	45.0	SASSIE RIDGE	PILLOW	4200	5/01/98	---	29.5S	59.4	24.1	
DISCOVERY BASIN	7050	4/30/98	29	11.2	17.9	10.0	SAVAGE PASS	PILLOW	6170	5/01/98	---	17.8	43.8	26.7	
DIX HILL	6400	5/03/98	0	.0	8.5	4.4	SAWMILL RIDGE		4700	5/02/98	56	25.8	64.4	28.2	
DOMMERIE FLATS	2200	4/29/98	0	.0	--	SCREIBERS MDW	AM	3400	5/01/98	---	48.0E	74.0	56.2		
EAST FORK R.S.	5400	4/25/98	0	.0	5.6	.9	SHEEP CANYON	PILLOW	4050	5/01/98	---	29.7S	34.7	34.7	
EAST RAGGED SADDLE	3740	5/03/98	0	.0	24.0	--	SKALKAHO	PILLOW	7260	5/01/98	---	21.6	42.6	26.2	
EASY PASS	AM 5200	5/01/98	---	93.0E	132.0	85.4	SKITWISI RIDGE		5110	5/05/98	23	11.2	48.7	30.3	
ELBOW LAKE	PILLOW	3200	5/01/98	---	15.7S	49.1	27.8	SKOOKUM CREEK	PILLOW	3920	5/01/98	---	7.5S	41.1	26.4
EMERY CREEK PILLOW	4350	5/01/98	---	.3	18.1	8.5	SLIDE ROCK MOUNTAIN		7100	4/25/98	39	13.6	24.6	17.2	
FATTY CREEK	5500	4/27/98	44	16.8	41.3	23.6	SPENCER MDW	PILLOW	3400	5/01/98	---	25.5S	46.1	17.2	
FISH CREEK	8000	4/24/98	49	14.7	17.2	12.4	SPIRIT LAKE	PILLOW	3100	5/01/98	---	.0S	.0	.3	
FISH LAKE	3370	4/28/98	46	19.6	45.6	22.4	SPOTTED BEAR MTN.		7000	5/01/98	0	.0	17.1	9.6	
FISH LAKE	PILLOW	3370	5/01/98	---	17.9S	49.0	25.0	STAHL PEAK	PILLOW	6030	5/01/98	---	33.2	51.9	36.5
FLATTOP MTN PILLOW	6300	5/01/98	---	36.0	66.9	48.4	STAMPEDE PASS	PILLOW	3860	5/01/98	---	33.8S	83.5	39.1	
FLEECER RIDGE	7500	5/01/98	19	6.2	21.4	8.4	STEMPLE PASS		6600	4/29/98	10	3.2	14.8	10.3	
FOURTH OF JULY SUM	3200	5/01/98	0	.0	.0	.0	STEVENS PASS	PILLOW	4070	5/01/98	---	22.1S	65.9	32.1	
FREEZEOUT CR. TRAIL	3500	4/28/98	7	3.9	13.7	7.0	STEVENS PASS	SAND SD	3700	5/01/98	37	17.2	58.0	28.7	
FROHNER MDWS PILLOW	6480	5/01/98	---	5.2	10.4	7.1	STORM LAKE		7780	4/30/98	45	15.0	22.4	15.0	
GRASS MOUNTAIN #2	2900	5/02/98	0	.0	.0	2.3	STRYKER BASIN		6180	4/27/98	70	29.7	47.2	35.8	
GRAVE CREEK PILLOW	4300	5/01/98	---	2.4	17.3	9.0	STUART MOUNTAIN		7400	4/27/98	67	27.6	53.6	32.3	
GREEN LAKE	PILLOW	6000	5/01/98	---	23.8S	33.9	SUNSET	PILLOW	5540	5/01/98	---	11.7	48.1	36.5	
GRIFFIN CR DIVIDE	5150	4/28/98	0	.0	13.6	6.3	SURPRISE LKS	PILLOW	4250	5/01/98	---	45.3S	74.2	36.1	
GROUSE CAMP	PILLOW	5380	5/01/98	---	12.0S	22.2	TEN MILE LOWER		6600	4/29/98	6	1.9	7.2	5.4	
HAND CREEK PILLOW	5030	5/01/98	---	.0	14.6	8.3	TEN MILE MIDDLE		6800	4/29/98	25	8.2	14.8	12.4	
HARTS PASS	PILLOW	6500	5/01/98	---	42.4S	59.6	42.0	THUNDER BASIN		4200	4/29/98	40	16.0	35.8	21.8
HELL ROARING DIVIDE	5770	4/30/98	46	19.6	42.1	30.1	TINKHAM CREEK	PILLOW	3000	5/01/98	---	15.0S	47.4	16.7	
HERRIG JUNCTION	4850	4/27/98	38	17.0	37.3	23.2	TOUCET #2	PILLOW	5530	5/01/98	---	18.3	56.6	27.3	
HIGH RIDGE	PILLOW	4980	5/01/98	---	7.8S	31.6	12.4	TRAPPING CK UP	CAN.	4100	5/03/98	0	.0	4.6	.3
HOLBROOK		4530	5/01/98	0	.0	9.9	1.7	TRINKUS LAKE		6100	5/01/98	57	27.4	69.6	43.1
HOODOO BASIN PILLOW	6050	5/01/98	---	31.5	76.7	47.2	TROUGH #2	PILLOW	5310	5/01/98	---	7.8S	2.5	--	
HUMBOLDT GLC PILLOW	4250	5/01/98	---	1.5	4.4	8.9	TRUMAN CREEK		4060	4/28/98	0	.0	2.3	.6	
HURRICANE		4500	4/27/98	35	15.2	23.6	TUNNEL AVENUE		2450	4/29/98	15	6.0	32.1	12.7	
INTERGAARD		6450	4/29/98	19	7.4	13.0	TV MOUNTAIN		6800	4/27/98	37	13.4	31.6	18.7	
JUNE LAKE	PILLOW	3200	5/01/98	---	21.6S	49.6	TWELVE MILE	PILLOW	5600	5/01/98	---	.2	26.4	12.4	
KRAFT CREEK PILLOW	4750	5/01/98	---	.0	22.3	5.8	TWIN CAMP		4100	5/02/98	30	12.1	38.6	--	
LESTER CREEK		3100	5/02/98	30	13.1	35.6	TWIN CREEKS		3580	5/01/98	0	.0	13.0	1.8	
LOGAN CREEK		4300	4/28/98	0	.0	7.3	TWIN LAKES	PILLOW	6400	5/01/98	---	29.3	69.4	39.8	
LOLO PASS	PILLOW	5240	5/01/98	---	14.0	55.0	TWIN SPIRIT DIVIDE		3480	5/03/98	0	.0	15.1	--	
LONE PINE	PILLOW	3800	5/01/98	---	35.3S	60.1	27.5	UPPER HOLLAND LAKE		6200	5/01/98	53	24.8	52.8	35.2
LOOKOUT	PILLOW	5140	5/01/98	---	18.7	47.8	29.3	UPPER WHEELER	PILLOW	4400	5/01/98	---	6.9S	12.0	4.8
LOST HORSE	PILLOW	5000	5/01/98	---	16.2S	20.5	8.2	WARM SPRINGS	PILLOW	7800	5/01/98	---	21.2	36.7	24.9
LOST LAKE	PILLOW	6110	5/01/98	---	38.2	102.6	63.0	WATSON LAKES	AM	4500	5/01/98	---	59.0E	81.0	67.2
LOWER SANDS CREEK #2															



U. S. DEPARTMENT OF AGRICULTURE
NATIONAL RESOURCES CONSERVATION SERVICE
WASHINGTON STATE OFFICE, SPOKANE

Spokane River Basin



*Based on selected stations

The May 1 forecasts for summer runoff within the Spokane River Basin are 50% of average near Post Falls and 57% of average at Long Lake. These forecasts dropped about 10% from last month. The forecast is based on a basin snowpack that is 50% of average and precipitation that is 83% of average for the water year. Precipitation for April was much below normal at 61% of average. Streamflow on the Spokane River at Long Lake, was 79% of average for April. May 1 storage in Coeur d'Alene Lake, was 181,500 acre feet, 74% of average, and 76% of capacity. Snowpack at Quartz Peak SNOTEL site contained 9.7 inches of water, compared to the average May 1 reading of 18.6 inches. Average temperatures in the Spokane Basin were 2 degrees above normal.

For more information contact your local Natural Resources Conservation Service office.

Spokane River Basin

Streamflow Forecasts - May 1, 1998

SPOKANE near Post Falls (2)	MAY-SEP	548	769		919	50		1069	1290	1846
	MAY-JUL	486	703		851	49		999	1216	1749
SPOKANE at Long Lake	MAY-JUL	691	918		1073	54		1228	1455	1975
	MAY-SEP	859	1092		1250	57		1408	1641	2198

SPOKANE RIVER BASIN

Reservoir Storage (1000 AF) - End of April

SPOKANE RIVER BASIN

Watershed Snowpack Analysis - May 1, 1998


Reservoir	Capacity	Usable Storage			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
COEUR D'ALENE	238.5	181.5	546.5	246.7	SPOKANE RIVER	10	32	50
					NEWMGN LAKE	1	34	52

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.



The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Quartz Peak SNOTEL Elevation 4700 ft.



Colville - Pend Oreille River Basins



*Based on selected stations

The forecast for the Kettle River streamflow is 105% of average; the Pend Oreille below Box Canyon, 59%; and the Priest River near the town of Priest River, 62% of average for the summer runoff period. April streamflow was 88% of average on the Pend Oreille River; 110% on the Columbia at the International Boundary; and 172% on the Kettle River. May 1 snow cover was 65% of average in the Pend Oreille Basin and 79% of average in the Kettle River Basin. Precipitation during April was 59% of average, bringing the year-to-date precipitation to 90% of average. Reservoir storage in Roosevelt and Banks Lakes was not available in time for this publication. Average temperatures were 3 degrees above normal.

Colville - Pend Oreille River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg. (1000AF)	
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
PEND OREILLE Lake Inflow (1,2)	MAY-JUL	4805	6149	6760	61	7371	8715	11070	
	MAY-SEP	5380	6872	7550	61	8228	9720	12290	
PRIEST nr Priest River (1,2)	MAY-SEP	260	370	420	62	470	580	680	
PEND OREILLE bl Box Canyon (1,2)	MAY-SEP	4675	6528	7370	59	8212	10065	12430	
	MAY-JUN	3517	4915	5550	59	6185	7583	9410	
CHAMOKANE CREEK near Long Lake	MAY-AUG	4.75	6.93	8.40	99	9.87	12.05	8.52	
COLVILLE at Kettle Falls	MAY-SEP	52	68	78	93	89	104	84	
	MAY-JUN	40	51	58	91	65	76	64	
KETTLE near Laurier	MAY-SEP	1394	1554	1662	105	1770	1930	1582	
	MAY-JUN	1172	1284	1360	104	1436	1548	1314	
COLUMBIA at Birchbank (1,2)	MAY-JUL	22932	25179	26200	82	27221	29468	32090	
	MAY-SEP	29436	32299	33600	82	34901	37764	40760	
	MAY-JUN	16264	17836	18550	82	19264	20836	22620	

COLVILLE - PEND OREILLE RIVER BASINS Reservoir Storage (1000 AF) - End of April

COLVILLE - PEND OREILLE RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	Usable Storage ***			Watershed	Number of Data Sites	This Year as % of Last Yr	Average
		This Year	Last Year	Avg				
ROOSEVELT	NO REPORT				COLVILLE RIVER	0	0	0
BANKS	NO REPORT				PEND OREILLE RIVER	90	40	65
					KETTLE RIVER	8	55	79

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.


The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Bunchgrass Meadows SNOTEL Elevation 5000 ft.



Okanogan - Methow River Basins



*Based on selected stations

Summer runoff forecast for the Okanogan River is 73% of average; the Similkameen River, 73%; the Methow River, 94%; and Salmon Creek, 122% of average. May 1 snow cover in the Okanogan was 82% of average; the Methow, 95%; and the Similkameen River, 57%. Salmon Meadows SNOTEL site above Conconully Lake had an May 1 reading of 436% of average. April precipitation in the Okanogan-Methow was 93% of average, with precipitation for the water year at 105% of average. April streamflow for the Methow River was 133% of average; 125% for the Okanogan River; and 81% for the Similkameen. Snow-water-content at the Salmon Meadows SNOTEL, near Conconully, was 4.8 inches. Average for this site is 1.1 inches on May 1. Combined storage in the Conconully Reservoirs was 23,800 acre feet, which is 101% of capacity and 149% of the May 1 average.

Okanogan - Methow River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
SIMILKAMEEN near Nighthawk (1)	MAY-JUL	610	803	890	74	977	1170	1205	
	MAY-SEP	663	860	950	73	1040	1237	1300	
	MAY-JUN	488	668	750	74	832	1012	1014	
OKANOGAN near Tonasket (1)	MAY-JUL	476	812	965	73	1118	1454	1328	
	MAY-SEP	548	914	1080	73	1246	1612	1484	
	MAY-JUN	381	669	800	73	931	1219	1094	
SALMON CREEK near Conconully	MAY-JUL	12.0	18.0	22	122	26	32	18.0	
	MAY-SEP	12.5	18.7	23	122	27	34	18.9	
METHOW RIVER near Pateros	MAY-SEP	693	757	800	94	843	907	854	
	MAY-JUL	647	704	742	94	780	837	786	
	MAY-JUN	537	590	626	95	662	715	659	

OKANOGAN - METHOW RIVER BASINS Reservoir Storage (1000 AF) - End of April

OKANOGAN - METHOW RIVER BASINS Watershed Snowpack Analysis - May 1, 1998

Reservoir	Capacity	Usable Storage			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
SALMON LAKE	10.5	10.3	8.7	8.0	OKANOGAN RIVER	22	60	82
CONCONULLY RESERVOIR	13.0	13.5	12.3	8.0	OMAK CREEK	1	89	167
					SANPOIL RIVER	0	0	0
					SIMILKAMEEN RIVER	5	44	57
					CONCONULLY LAKE	1	53	436
					METHOW RIVER	3	62	95


* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.

The average is computed for the 1961-1990 base period.


(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.

(2) - The value is natural flow - actual flow may be affected by upstream water management.

Salmon Meadows SNOTEL Elevation 4500 ft.



Wenatchee - Chelan River Basins



*Based on selected stations

Precipitation during April was 64% of average in the basin and 104% for the year-to-date. Runoff for the Entiat River is forecast to be 101% of average for the summer. The May-September forecast for the Chelan River is for normal flows; for the Wenatchee River at Plain it is 93%; and for the Stehekin it is 100% of average. Icicle, Stemilt and Squilchuck Creeks are all expected to have near normal flows this summer as well. April streamflow on the Chelan River was 108% of average, and the Wenatchee River averaged normal flows. May 1 snowpack in the Wenatchee Basin was 90% of average. The Chelan Basin was 100% of average; Colockum Ridge was 312%; and Stemilt Creek was 144% of average. Snowpack in the Entiat River Basin was near normal. Reservoir storage in Lake Chelan was 312,800 acre feet, or 70% of May 1 average and 46% of capacity. Lyman Lake SNOTEL had the most snow water with 68.5 inches of water. This site would normally have 58.7 inches on May 1. Temperatures were slightly above normal for April.

For more information contact your local Natural Resources Conservation Service office.

Wenatchee - Chelan River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		Drier			Wetter				
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
CHELAN RIVER near Chelan	MAY-SEP	929	995	1040	100	1085	1151	1041	
	MAY-JUL	812	873	915	101	957	1018	905	
	MAY-JUN	589	655	700	101	745	811	693	
STEHEKIN near STEHEKIN	MAY-SEP	672	718	750	100	782	828	751	
	MAY-JUL	548	594	625	100	656	702	625	
	MAY-JUN	378	427	460	100	493	542	462	
ENTIAT RIVER near Ardenvoir	MAY-SEP	189	202	211	101	220	233	208	
	MAY-JUL	169	182	191	102	200	213	188	
	MAY-JUN	132	145	155	103	165	178	150	
WENATCHEE at Plain	MAY-SEP	845	919	970	93	1021	1095	1042	
	MAY-JUL	761	819	858	93	897	955	925	
	MAY-JUN	568	613	644	90	675	720	716	
STEMILT nr Wenatchee (miners in)	MAY-SEP	85	112	130	94	148	175	138	
ICICLE CREEK near Leavenworth	MAY-SEP	268	278	285	93	292	302	305	
	MAY-JUL	236	250	260	93	270	284	279	
	MAY-JUN	166	192	210	94	228	254	224	

WENATCHEE - CHELAN RIVER BASINS Reservoir Storage (1000 AF) - End of April

WENATCHEE - CHELAN RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	Usable Storage			Watershed	Number of Data Sites	This Year as % of Last Yr Average	
		This Year	Last Year	Avg			Last Yr	Average
CHELAN LAKE	676.1	312.8	233.6	448.8	CHELAN LAKE BASIN	4	63	100
					ENTIAT RIVER	1	41	606
					WENATCHEE RIVER	9	48	90
					SQUILCHUCK CREEK	0	0	0
					STEMILT CREEK	1	58	144
					COLOCKUM CREEK	1	116	312

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.


The average is computed for the 1961-1990 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural flow - actual flow may be affected by upstream water management.

Pope Ridge SNOTEL
Elevation 3540 ft.



Yakima River Basin



*Based on selected stations

May 1 reservoir storage for the five major reservoirs was 958,400 acre feet, or 122% of average. May 1 summer streamflow forecasts are for near to slightly below normal in the Yakima Basin. Forecasts for the Yakima River at Cle Elum, are for 86% of average; Naches River, 95%; the Yakima River near Parker, 88%; Ahtanum Creek, 92%; and the Tieton River, 98%. The Klickitat River near Glenwood is forecast to have 111% of average flows this summer. Volume forecasts for the Yakima Basin are for natural flow. As such, they may differ from the U.S. Bureau of Reclamation's forecast for the total water supply available, which includes irrigation return flow. April streamflows within the basin were: the Yakima River near Kiona, 103% of average; the Yakima River near Cle Elum, 91%; and the Naches River at 111%. May 1 snowpack was 106% based upon 15 snow courses and SNOTEL readings within the Yakima Basin. Precipitation was 58% of average for April and 102% for the water year-to-date.

Yakima River Basin

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
KEECHELUS LAKE INFLOW	MAY-JUL	67	77	84	87	90	100	96	
	MAY-SEP	72	85	93	87	101	114	107	
	MAY-JUN	53	64	71	87	77	88	81	
KACHESS LAKE INFLOW	MAY-JUL	62	68	73	85	78	85	86	
	MAY-SEP	64	73	78	85	84	92	92	
	MAY-JUN	52	59	63	85	67	74	74	
CLE ELUM LAKE INFLOW	MAY-JUL	257	275	288	85	301	319	339	
	MAY-SEP	282	305	320	85	335	358	378	
	MAY-JUN	201	221	235	85	249	269	276	
YAKIMA at Cle Elum	MAY-JUN	406	444	470	86	496	534	546	
	MAY-JUL	496	537	565	86	593	634	657	
	MAY-SEP	548	600	635	86	670	722	740	
BUMPING LAKE INFLOW	MAY-SEP	92	103	110	94	117	128	117	
	MAY-JUL	87	95	101	95	107	115	106	
	MAY-JUN	73	80	85	99	90	97	86	
AMERICAN RIVER near Nile	MAY-SEP	95	104	110	108	116	125	102	
	MAY-JUL	86	94	100	109	105	114	92	
	MAY-JUN	67	75	81	108	87	95	75	
RIMROCK LAKE INFLOW	MAY-SEP	174	189	200	98	211	226	204	
	MAY-JUL	146	157	165	99	173	184	167	
	MAY-JUN	101	116	127	98	138	153	129	
NACHES near Naches	MAY-SEP	561	614	650	95	686	739	686	
	MAY-JUL	512	555	585	96	615	658	609	
	MAY-JUN	419	458	485	96	512	551	505	
AHTANUM CREEK nr Tampico (2)	MAY-SEP	27	32	35	92	38	44	38	
	MAY-JUL	24	29	32	93	35	39	34	
	MAY-JUN	19.7	23	26	93	29	32	28	
YAKIMA near Parker	MAY-SEP	1212	1315	1385	88	1455	1558	1580	
	MAY-JUL	1065	1157	1220	88	1283	1375	1390	
	MAY-SEP	1212	1315	1385	88	1455	1558	1580	
KLICKITAT near Glenwood	MAY-JUN	81	90	96	110	102	110	87	
	MAY-SEP	111	122	130	111	138	149	117	

YAKIMA RIVER BASIN					YAKIMA RIVER BASIN			
Reservoir Storage (1000 AF) - End of April					Watershed Snowpack Analysis - May 1, 1998			
Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of Last Yr Average	
	Year	This Year	Last Year	Avg				
KEECHELUS	157.8	150.2	118.9	119.0	YAKIMA RIVER	15	54	106
KACHESS	239.0	214.9	185.9	197.0	AHTANUM CREEK	2	74	143
CLE ELUM	436.9	407.0	279.1	308.0				
BUMPING LAKE	33.7	21.2	15.1	15.0				
RIMROCK	198.0	165.1	119.0	144.0				



* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.

The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.

(2) - The value is natural flow - actual flow may be affected by upstream water management.

Walla Walla River Basin



*Based on selected stations

April precipitation was 63% of average, bringing the year-to-date precipitation to 80% of average. May 1 snowpack was 66% of average. The summer forecast is for 77% of average streamflow in the Snake River below Lower Granite Dam, 93% for the Grande Ronde at Troy, and 90% for Mill Creek. April streamflow was 115% of average for the Walla Walla River; 100% for the Snake River below Lower Granite Dam; and 77% for the Grande Ronde River near Troy. The Touchet SNOTEL site had 7.8 inches of snow-water-equivalent. The average May 1 reading for this site is 2.5 inches. Average temperatures were near normal for the area.

Walla Walla River Basin

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		<===== Drier =====		Chance Of Exceeding *			Wetter =====>		
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
GRANDE RONDE at Troy (1)	MAY-JUL	566	734	810	93	886	1054	872	
	MAY-SEP	628	815	900	93	985	1172	970	
SNAKE blw Lower Granite Dam (1,2)	MAY-JUL	9510	11772	12800	76	13828	16090	16940	
	MAY-SEP	11295	13912	15100	77	16288	18905	19650	
MILL CREEK at Walla Walla	MAY-SEP	3.57	5.46	6.75	90	8.04	9.93	7.50	
	MAY-JUL	3.43	5.32	6.60	90	7.88	9.77	7.30	
	MAY-JUN	3.39	5.18	6.40	90	7.62	9.41	7.10	
SF WALLA WALLA near Milton-Freewater	MAY-JUL	25	29	32	85	34	38	37	
	MAY-SEP	36	41	44	88	47	52	50	

WALLA WALLA RIVER BASIN

Reservoir Storage (1000 AF) - End of April

WALLA WALLA RIVER BASIN

Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of Last Yr Average	
		This Year	Last Year	Avg				
					WALLA WALLA RIVER	2	30	66

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.


The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Touchet #2 SNOTEL Elevation 5530 ft.



Cowlitz - Lewis River Basins



*Based on selected stations

The forecast for summer runoff in the Lewis River Basin is for 89% of average. The Cowlitz River at Castle Rock, is forecast for 96% of average runoff. April streamflow was 67% of average for the Cowlitz River, and 68% of average for the Lewis River. April precipitation was 50% of average. It was 105% of average for the water-year. May 1 snow cover for the Cowlitz River was 106%, and the Lewis River was 129% of average. The Paradise Park SNOTEL recorded the most water-content for the basin with 70 inches of water. Average May 1 water-content is 61.8 inches. Average temperatures were 1 to 2 degrees above normal during April.

For more information contact your local Natural Resources Conservation Service office.

Cowlitz - Lewis River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						
		<===== Drier =====		===== Wetter =====>				
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
LEWIS at Ariel (2)	MAY-JUL	496	570	620	89	670	744	697
	MAY-SEP	633	709	760	89	811	887	850
	MAY-JUN	402	469	515	89	561	628	579
COWLITZ R. bl Mayfield Dam (2)	MAY-SEP	630	1094	1410	92	1726	2190	1531
	MAY-JUL	532	922	1188	92	1454	1844	1292
	MAY-JUN	424	740	955	92	1170	1486	1038
COWLITZ R. at Castle Rock (2)	MAY-SEP	933	1527	1930	96	2333	2927	2021
	MAY-JUL	770	1264	1600	95	1936	2430	1679
	MAY-JUN	610	1009	1280	95	1551	1950	1349
KLICKITAT near Glenwood	MAY-JUN	81	90	96	110	102	110	87
	MAY-SEP	111	122	130	111	138	149	117

COWLITZ - LEWIS RIVER BASINS Reservoir Storage (1000 AF) - End of April

COWLITZ - LEWIS RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	Usable Storage ***			Watershed	Number of Data Sites	This Year as % of Last Yr	This Year as % of Average
		This Year	Last Year	Avg				
					LEWIS RIVER	4	56	129
					COWLITZ RIVER	7	64	106

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.


The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Paradise SNOTEL Elevation 5120 ft.



White - Green River Basins



*Based on selected stations

Summer runoff is forecast to be 68% of average for the Green River. A 10% drop from last month. The White River should see near to slightly below normal flows while the Nisqually River will most likely experience below normal flows this summer. May 1 snowpack was 123% of average in the White River Basin; and 78% in the Green River Basin. Water-content on May 1 at the Morse Lake SNOTEL, at an elevation of 5,400 feet, was 68.3 inches. This site has an May 1 average of 44.4 inches. April precipitation was 50% of average, bringing the water year-to-date to 93% of average for the basins. April temperatures were slightly above average.

For more information contact your local Natural Resources Conservation Service office.

White - Green River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg. (1000AF)	
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)		
GREEN RIVER below Howard Hanson Dam	MAY-JUL	79	101	116	68	131	153	170	
	MAY-SEP	94	118	135	68	152	176	198	
	MAY-JUN	68	87	100	68	113	132	147	

WHITE - GREEN RIVER BASINS Reservoir Storage (1000 AF) - End of April

WHITE - GREEN RIVER BASINS Watershed Snowpack Analysis - May 1, 1998

Reservoir	Usable Capacity	Usable Storage ***			Watershed	Number of Data Sites	This Year as % of Last Yr	Average
		This Year	Last Year	Avg				
					WHITE RIVER	3	76	123
					GREEN RIVER	6	34	78


* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.

The average is computed for the 1961-1990 base period.



(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.

(2) - The value is natural flow - actual flow may be affected by upstream water management.

Stampede Pass SNOTEL Elevation 3860 ft.



Central Puget Sound River Basins



*Based on selected stations

Forecast for spring and summer flows are: 77% for the Cedar River near Cedar Falls; 73% for the Rex River; 90% for the South Fork of the Tolt River; and 75% for the Cedar River at Cedar Falls. The Cedar River at Cedar Falls stream gage may be affected by upstream reservoir control. Basin-wide precipitation for April was 47% of average, bringing the water-year-to-date to 92% of average. May 1 snow cover in the Cedar River Basin was 48%; the Tolt River Basin was 28%; the Snoqualmie River Basin was 78%; and the Skykomish River Basin was 65% of average. Stevens Pass SNOTEL, at 4,070 feet, had 22.1 inches of water content. Average May 1 water content is 32.1 inches. April temperatures were 1 to 2 degrees above normal.

For more information contact your local Natural Resources Conservation Service office.

Central Puget Sound River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>		
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	(% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
CEDAR near Cedar Falls	MAY-JUL	30	37	42	75	47	54	56
	MAY-SEP	35	43	49	77	54	62	64
	MAY-JUN	28	33	37	77	40	46	47
REX near Cedar Falls	MAY-JUL	8.5	11.8	14.0	73	16.2	19.5	19.2
	MAY-SEP	10.0	13.7	16.2	73	18.7	22	22
	MAY-JUN	7.6	10.2	12.0	73	13.8	16.4	16.5
CEDAR RIVER at Cedar Falls	MAY-JUL	8.9	27	40	74	53	71	54
	MAY-SEP	4.9	26	41	75	56	77	55
	MAY-JUN	15.9	29	39	74	48	61	52
SOUTH FORK TOLT near Index	MAY-JUL	7.4	8.9	9.9	87	10.9	12.4	11.4
	MAY-SEP	9.3	11.2	12.5	90	13.8	15.7	13.9
	MAY-JUN	6.11	7.32	8.14	88	8.96	10.17	9.30

CENTRAL PUGET SOUND RIVER BASINS Reservoir Storage (1000 AF) - End of April

CENTRAL PUGET SOUND RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity			Watershed	Number of Data Sites	This Year as % of	
	This Year	Last Year	Avg			Last Yr	Average
				CEDAR RIVER	4	24	48
				TOLT RIVER	1	18	28
				SNOQUALMIE RIVER	3	41	78
				SKYKOMISH RIVER	2	32	65

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.



The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Stevens Pass SNOTEL Elevation 4070 ft.



North Puget Sound River Basins



*Based on selected stations

Forecast for the Skagit River streamflow is for 85% of average for the spring and summer period. April streamflow in the Skagit River was 79% of average. Other forecast points included the Baker River at 86%; and Thunder Creek at 90% of average. Basin-wide precipitation for April was only 47% of average, bringing water-year-to-date to 90% of average. May 1 snow cover in the Skagit River Basin was 85%; the Baker River Basin was 93%; and the Nooksack River Basin dropped to 56% of average. Rainy Pass SNOTEL, at 4,780 feet, had 29 inches of water content. Average May 1 water content is 36.8 inches. May 1 Skagit River reservoir storage was 112% of average and 51% of capacity. Average April temperatures were about 2 degrees above normal for the basin.

North Puget Sound River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		<===== Drier =====		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	% AVG. (%)	30% (1000AF)	10% (1000AF)		
THUNDER CREEK near Newhalem	MAY-JUL	162	177	188	90	199	214	209	
	MAY-SEP	252	268	278	90	288	304	308	
	MAY-JUN	86	104	116	90	128	146	129	
SKAGIT near Newhalem (2)	MAY-JUL	1255	1341	1400	85	1459	1545	1649	
	MAY-SEP	1513	1607	1670	85	1733	1827	1961	
	MAY-JUN	879	975	1040	85	1105	1201	1224	
BAKER RIVER near Concrete	MAY-JUL	524	575	609	87	643	694	703	
	MAY-SEP	675	750	802	86	854	929	930	
	MAY-JUN	352	400	433	91	466	514	478	

NORTH PUGET SOUND RIVER BASINS Reservoir Storage (1000 AF) - End of April

NORTH PUGET SOUND RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
ROSS	1404.1	719.5	695.4	644.4	SKAGIT RIVER	13	56	85
DIABLO RESERVOIR	90.6	87.8	87.4	---	BAKER RIVER	4	70	93
GORGE RESERVOIR	NO REPORT				NOOKSACK RIVER	2	42	56

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.



The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Rainy Pass SNOTEL Elevation 4780 ft.



Olympic Peninsula River Basins



*Based on selected stations

May forecasts of runoff for streamflow in the Dungeness River Basin are 97% of average and 94% of average for the Elwha River. The Big Quilcene and Wynoochee rivers can expect near to above average runoff this summer. April precipitation was only 65% of average. Precipitation accumulated at 110% of average for the water year. April precipitation at Quillayute was 3.15 inches. The thirty-year average for May 1 is 7.15 inches. May 1 snow cover in the Olympic Basin was at 104% of average. The Mount Crag SNOTEL near Quilcene had 35.2 inches of snow-water-equivalent on May 1. Average for this site is 22.4 inches. Temperatures were near average for the month.

Olympic Peninsula River Basins

Streamflow Forecasts - May 1, 1998

Forecast Point	Forecast Period	Future Conditions						30-Yr Avg.	
		<<===== Drier =====>>		Chance Of Exceeding *		Wetter =====>			
		90% (1000AF)	70% (1000AF)	50% (Most Probable) (1000AF)	% AVG.	30% (1000AF)	10% (1000AF)		
DUNGENESS near Sequim	MAY-SEP	116	124	130	97	136	144	134	
	MAY-JUL	93	99	103	96	107	113	107	
	MAY-JUN	63	70	75	98	79	86	76	
ELWHA near Port Angeles	MAY-SEP	363	390	408	94	426	453	434	
	MAY-JUL	291	313	327	94	341	363	348	

OLYMPIC PENINSULA RIVER BASINS Reservoir Storage (1000 AF) - End of April

OLYMPIC PENINSULA RIVER BASINS Watershed Snowpack Analysis - May 1, 1998


Reservoir	Usable Capacity	*** Usable Storage ***			Watershed	Number of Data Sites	This Year as % of	
		This Year	Last Year	Avg			Last Yr	Average
					ELWHA RIVER	1	64	69
					MORSE CREEK	1	77	97
					DUNGENESS RIVER	1	128	94
					QUILCENE RIVER	1	124	157
					WYNOCHEE RIVER	0	0	0

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual flow will exceed the volumes in the table.

The average is computed for the 1961-1990 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
 (2) - The value is natural flow - actual flow may be affected by upstream water management.

Mount Crag SNOTEL Elevation 4050 ft.





Natural Resources Conservation Service

Washington State
Snow, Water and Climate Services

Program Contacts

Leonard Jordan
State Conservationist
W. 316 Boone Ave., Suite 450
Spokane, WA 99201-2348
phone: 509-323-2900
fax: 509-323-2909
ljordan@wa.nrcs.usda.gov

Chris Bieker
Public Affairs Specialist
W. 316 Boone Ave., Suite 450
Spokane, WA 99201-2348
phone: 509-323-2912
fax: 509-323-2909
cbieker@wa.nrcs.usda.gov

Scott Pattee
Water Supply Specialist
2021 E. College Way, Suite 214
Mount Vernon, WA 98273-2873
phone: 360-428-7684
fax: 360-424-6172
spattee@wa.nrcs.usda.gov

Chris Pacheco
Resource Conservationist
National Water and Climate Center
101 SW Main St., Suite 1600
Portland, OR 97204-3224
phone: 503-414-3056
fax: 503-414-3101
cpacheco@wcc.nrcs.usda.gov

Helpful Internet Addresses

NRCS Snow Survey and Climate Services Homepages

Washington:
<http://wcp.wsu.edu/nrcs/CoopSnoSrvy.htm>

Oregon:
<http://crystal.or.nrcs.usda.gov/snowsurveys/>

Idaho:
<http://id.nrcs.usda.gov/snow/snow.htm>

National Water and Climate Center (NWCC):
<http://www.wcc.nrcs.usda.gov/>

NWCC Anonymous FTP Server:
<ftp://wcc.nrcs.usda.gov>

USDA-NRCS Agency Homepages

Washington:
<http://wcp.wsu.edu/nrcs/>

NRCS National:
<http://www.ftw.nrcs.usda.gov/>



Natural Resources Conservation Service

Washington State
Snow, Water and Climate Services

Field Office Contacts

Eastern Washington

Jimmie Gleaton
District Conservationist
232 Williams Lake Road
Colville, WA 99114-9638
509-684-0937
jgleaton@wacolville.fsc.usda.gov

Randy Kelley
District Conservationist
1251 Second Ave. South, Room 101
Okanogan, WA 98840-9723
509-422-2750
rkelley@waokanogan.fsc.usda.gov

Andrea Mann-Lower
Wetland Specialist
215 Melody Lane, Rm #238
Wenatchee, WA 98801-9533
509-664-2717

David Chain
District Conservationist
607 E. Mountain View
Ellensburg, WA 98926
509-925-8580
dchain@waellensbu.fsc.usda.gov

Western Washington

John Gillies
District Conservationist
6975 Hannegen Road
Lynden, WA 98205-1535
360-354-2035
jgillies@walynden.fsc.usda.gov

Monica Hoover
Wetland Specialist
6128 Capitol Blvd. SE
Olympia, WA 98501-5271
360-753-9448
mhoover@wa.nrcs.usda.gov

Kerry Perkins
District Conservationist
111 East 3rd, Room 2B
Port Angeles, WA 98362-3009
360-457-5091
kperkins@wa.nrcs.usda.gov

Scott Pattee
Water Supply Specialist
2021 E. College Way, Suite 214
Mount Vernon, WA 98273-2873
360-428-7684
spattee@wa.nrcs.usda.gov

Data Collection Offices

Jon Lea
Oregon Data Collection Office
101 SW Main St., Suite 1300
Portland, OR 97204
503-414-3267
jlea@or.nrcs.usda.gov

Phil Morrissey
Idaho Data Collection Office
9173 West Barnes, Suite C
Boise, ID 83709
208-378-5741
pmorris@id.nrcs.usda.gov

Issued by

Pearlie S. Reed
Chief
Natural Resources Conservation Service
U.S. Department of Agriculture

Released by

Leonard Jordan
State Conservationist
Natural Resources Conservation Service
Spokane, Washington

The Following Organizations Cooperate with the Natural Resources Conservation Service in Snow Survey Work*:

Canada	Ministry of the Environment Investigations Branch, Victoria, British Columbia
State	Washington State Department of Ecology Washington State Department of Natural Resources
Federal	Department of the Army Corps of Engineers U.S. Department of Agriculture Forest Service U.S. Department of Commerce NOAA, National Weather Service U.S. Department of Interior Bonneville Power Administration Bureau of Reclamation Geological Survey National Park Service Bureau of Indian Affairs
Local	City of Tacoma City of Seattle Chelan County P.U.D. Pacific Power and Light Company Puget Sound Power and Light Company Washington Water Power Company Snohomish County P.U.D. Colville Confederated Tribes Spokane County Yakama Indian Nation Whatcom County Pierce County
Private	Okanogan Irrigation District Wenatchee Heights Irrigation District Newman Lake Homeowners Association Whitestone Reclamation District



★ ★ ★
175 01.010 PB8615738
6051 \$ 01.010 MAY 08 98
2286 MOUNT VERNON WA 98273

Rock Pointe Tower II, Suite 450
W. 316 Boone Avenue
Spokane, WA 99201-2349

FOR OFFICIAL USE ONLY

U. S. DEPT. OF AGRICULTURE
NATIONAL AGRICUL. LIBRARY
CURRENT SERIAL RECORDS
ROOM 002
BELTSVILLE, MD 20705-2351

**Washington
Basin Outlook Report**
Natural Resources Conservation Service
Spokane, WA

